

Genetic specificity of a plant–insect food web: Implications for linking genetic variation to network complexity

Matthew A. Barbour^{a,1}, Miguel A. Fortuna^b, Jordi Bascompte^b, Joshua R. Nicholson^a, Riitta Julkunen-Tiitto^c, Erik S. Jules^d, and Gregory M. Crutsinger^a

^aDepartment of Zoology, University of British Columbia, Vancouver, BC, Canada V6T 1Z4; ^bDepartment of Evolutionary Biology and Environmental Studies, University of Zurich, 8057 Zurich, Switzerland; ^cDepartment of Biology, University of Eastern Finland, FI-80101 Joensuu, Finland; and ^dDepartment of Biological Sciences, Humboldt State University, Arcata, CA 95521

Edited by Daniel S. Simberloff, The University of Tennessee, Knoxville, TN, and approved January 7, 2016 (received for review July 13, 2015)

Theory predicts that intraspecific genetic variation can increase the complexity of an ecological network. To date, however, we are lacking empirical knowledge of the extent to which genetic variation determines the assembly of ecological networks, as well as how the gain or loss of genetic variation will affect network structure. To address this knowledge gap, we used a common garden experiment to quantify the extent to which heritable trait variation in a host plant determines the assembly of its associated insect food web (network of trophic interactions). We then used a resampling procedure to simulate the additive effects of genetic variation on overall food-web complexity. We found that trait variation among host-plant genotypes was associated with resistance to insect herbivores, which indirectly affected interactions between herbivores and their insect parasitoids. Direct and indirect genetic effects resulted in distinct compositions of trophic interactions associated with each host-plant genotype. Moreover, our simulations suggest that food-web complexity would increase by 20% over the range of genetic variation in the experimental population of host plants. Taken together, our results indicate that intraspecific genetic variation can play a key role in structuring ecological networks, which may in turn affect network persistence.

species interactions | ecological networks | evolutionary ecology | community genetics

Network theory has provided both a conceptual and a quantitative approach for mapping interactions between species and making predictions about how the gain or loss of species will affect the structure and dynamics of ecological networks (1–3). Representing a network at the species level, however, makes the implicit assumption that each species consists of a homogenous population of individuals, all of which interact equally with individuals of different species. However, most populations are heterogeneous mixtures of individuals that vary in their phenotypes, and there is growing evidence that this intraspecific variation is an important factor governing the assembly of ecological communities (4–6). Consequently, there is a clear need to account for the role of intraspecific variation in structuring ecological networks (7).

Genetic variation is a key driver of intraspecific variation and many studies have now demonstrated direct and indirect genetic effects on species interactions (8–10) and the composition of communities across multiple trophic levels (11–14). This prior work forms a clear expectation that intraspecific genetic variation is capable of scaling up to affect the structure of an ecological network. In particular, we expect that network structure will be affected by genetic variation through at least two different mechanisms. For a food web (network of trophic interactions), genetic variation in the quality of a basal resource may alter the (*i*) abundances or (*ii*) phenotypes of consumer species or both (15). These direct genetic effects on consumers may then have cascading effects on the strength of trophic interactions between consumers and their predators (15), resulting in distinct compositions

of trophic interactions associated with different genotypes of the basal resource (Fig. 1). If such genetic specificity in the composition of trophic interactions occurs, then theory predicts that increasing genetic variation will result in more interactions per species (6, 16) and therefore greater food-web complexity (Fig. 2). Moreover, greater complexity may in turn affect food-web dynamics, as more complex food webs are predicted to be more robust to species extinctions (3, 17). However, whether genetic variation is capable of scaling up to affect food-web complexity is currently unclear.

In this study, we quantify the genetic specificity of trophic interactions and use these data to simulate the additive effects of genetic variation on food-web complexity. To do this, we used a common garden experiment of a host plant (26 genotypes of coastal willow, *Salix hookeriana*) and its associated food web of insect galls and parasitoids (Fig. 1). We focused on this plant–insect food web for three reasons. First, we have demonstrated in previous work that *S. hookeriana* (hereafter, willow) displays heritable variation in traits associated with leaf quality (36 traits, mean $H^2 = 0.72$) and plant architecture (4 traits, mean $H^2 = 0.27$), some of which are also associated with resistance to its community of galling herbivores (18). Second, the unique biology of galling insects makes them ideal for building quantitative food webs. In particular, galls provide a refuge

Significance

We know that the gain or loss of species can have cascading effects on food-web complexity; however, it is less clear whether the gain or loss of genetic variation within species, an often overlooked component of biodiversity, will similarly affect food-web structure. Here, we empirically identify how genetic variation within a host plant directly and indirectly affects its associated insect food web, resulting in distinct trophic interactions occurring on each host-plant genotype. Moreover, simulations of our empirical data suggest that higher levels of host-plant genetic variation lead to a more complex plant–insect food web. Our results suggest that preserving genetic variation within key species may be critical for maintaining complex and robust food webs under future environmental change.

Author contributions: M.A.B., E.S.J., and G.M.C. designed research; M.A.B. and J.R.N. performed research; M.A.F., J.B., and R.J.-T. contributed new reagents/analytic tools; M.A.B. analyzed data; and M.A.B., M.A.F., J.B., and G.M.C. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Data deposition: The data reported in this paper are available at the Dryad Digital Repository: [dx.doi.org/10.5061/dryad.g7805](https://doi.org/10.5061/dryad.g7805). Code to analyze the data is available at the GitHub Repository: [dx.doi.org/10.5281/zenodo.45210](https://doi.org/10.5281/zenodo.45210).

¹To whom correspondence should be addressed. Email: barbour@zoology.ubc.ca.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1513633113/-DCSupplemental.




Fig. 1. Genetic specificity of trophic interactions in a plant–insect food web. The species comprising the food web in this study include a host plant (coastal willow, *S. hookeriana*), four herbivorous gall-forming insects, and six insect parasitoids (species details in *Materials and Methods*). The plant–insect food web consists of 16 trophic interactions (4 willow–gall and 12 gall–parasitoid) aggregated from all plant individuals sampled in this common garden experiment, whereas each genotype subweb represents the trophic interactions aggregated from all plant individuals of the corresponding genotype. We depicted three genotype subwebs (of 26) to illustrate the differences in trophic interactions associated with each willow genotype. The width of each gray segment is proportional to the number of individuals associated with each trophic interaction. Note that we scaled the width of trophic interactions to be comparable among genotype subwebs, but not between subwebs and the aggregated food web, to emphasize the differences among subwebs.

for larva from attack by most generalist predators (19); therefore, galls and their natural enemies often form a distinct subset of the larger food web associated with host plants. In our system, all of the natural enemies are insect parasitoids that complete their development within the gall after parasitizing larva, making it easy to identify and quantify all of the trophic interactions within this food web. Third, the biology of galls is also ideal for identifying the mechanisms mediating trophic interactions. In particular, gall size is a key trait that affects the ability of parasitoids to successfully oviposit through the gall wall and into the larva within the gall (i.e., larger galls provide a refuge from parasitism) (20). Moreover, gall size is determined, in part, by the genotype of the plant (20), so we have a clear mechanism by which genetic variation can affect the strength of trophic interactions. Taken together, our study seeks to examine how intraspecific genetic variation influences the structure of ecological networks. In doing so, our study takes a crucial step toward a more predictive understanding of how the gain or loss of genetic variation will affect the dynamics of ecological networks.

Results and Discussion

Quantifying the Genetic Specificity of the Plant–Insect Food Web. In concordance with previous work in this system (18), we observed clear differences in the abundance of three of the four gall-forming insects among willow genotypes [multivariate generalized linear model (GLM), $\chi^2_{25,119} = 202.40$, $P = 0.001$] (*SI Appendix, Table S1*).

Specifically, we found that the average abundance of leaf, bud, and apical-stem galls varied 10-, 8-, and 1.4-fold among willow genotypes, respectively (Fig. 3 *A–C*). This variation resulted in 69% dissimilarity in the average composition of galls among willow genotypes ($F_{22,89} = 1.96$, $P = 0.001$). Moreover, we found that the average diameter of leaf galls varied twofold among willow genotypes (Fig. 3*D*). This observed genetic specificity in the abundance and phenotypes of insect herbivores corroborates decades of work in other plant–gall (8, 11, 20) and plant–herbivore systems (12, 21).

Importantly, however, our extensive screening of willow phenotypes (*Materials and Methods*) enabled us to identify traits that may be mediating the genetic specificity of trophic interactions with gall-forming insects. In particular, we found that leaf carbon-to-nitrogen ratio (C:N), certain leaf secondary metabolites (flavonones/flavanonols PC1), and plant size were associated with changes in the abundance of gall-forming insects (multivariate GLM, $\chi^2_{3,104} = 28.44$, $P = 0.004$) (*SI Appendix, Table S2*), whereas leaf gall diameter was associated with variation in a different suite of leaf secondary metabolites (salicylates/tannins PC1 and flavones/flavonols PC1; weighted linear model, $F_{2,59} = 8.27$, $P < 0.001$) (*SI Appendix, Table S2*). These results highlight that accounting for intraspecific variation in multiple plant traits is important for predicting antagonistic interactions between plants and insect herbivores (18) and should therefore be incorporated into mechanistic models of food-web structure.

We found that the effects of willow genetic variation extended beyond pairwise interactions with herbivores (11, 12, 21) and simple tritrophic interactions (8–10, 20) to determine the assembly of the network of gall–parasitoid interactions (multivariate GLM, $\chi^2_{25,119} = 357.10$, $P = 0.001$) (*SI Appendix, Table S1*). In particular, we found that the frequency of parasitism from three parasitoids (*Platygaster* sp., *Mesopolobus* sp., and *Torymus* sp.) on leaf galls varied 270%, 30%, and 40% among willow genotypes, respectively (Fig. 4 *A–C*). This variation resulted in




Fig. 2. Conceptual model of how increasing genetic variation (number of shades of green circles) results in greater food-web complexity (number of interactions per species). If different genotypes of a basal resource are associated with distinct compositions of trophic interactions (i.e., genetic specificity of trophic interactions), then increasing genetic variation in the resource will result in a more complex food web because of the increase in the number of interactions per species at all three trophic levels. Colors correspond to different trophic levels (green, basal resource; blue, primary consumer; orange, secondary consumer), whereas different shapes within each trophic level correspond to different species.

natural selection can play an important role in shaping consumer-resource dynamics (22, 23). Understanding how evolutionary processes affect the structure and dynamics of ecological networks, and vice versa (24, 25), is likely a fruitful topic for future research.

Simulating the Additive Effects of Genetic Variation on Network Complexity. To examine this, we used our empirical data to simulate how the complexity of the plant–insect food web would change across different levels of willow genetic variation (*Materials and Methods*). After accounting for sampling effort (Fig. 6, dashed line), our simulations suggest that food-web complexity would increase by 20% with increasing genetic variation (Fig. 6). This positive relationship was primarily due to an increased likelihood of sampling genotypes with complementary trophic interactions, as we found that willow genotypes differed by 73% in the average composition of their trophic interactions (Fig. 6, *Inset*). To more precisely understand the relationship between genetic variation, the addition of complementary interactions,




Fig. 5. Variation in the size and abundance of leaf galls on willows is associated with changes in the strength and composition of gall–parasitoid interactions. (A and B) In general, the proportion of leaf galls parasitized by both *Platygaster* (blue, solid line) and *Mesopolobus* (green, short-dashed line) decreases as gall size increases, whereas *Torymus* (orange, long-dashed line) exhibits the opposite pattern. On willows with small leaf galls (<8 mm), however, *Mesopolobus* had the highest attack rate at low gall abundances (1–4 leaf galls per branch; $n = 46$ per parasitoid species), whereas *Platygaster* was the dominant parasitoid at high gall abundances (5–22 leaf galls per branch; $n = 35$ per parasitoid species). Lines correspond to slopes estimated from GLMs. Points were jittered slightly to avoid overlapping values.




Fig. 6. Simulations of our empirical data indicate that increasing willow (*S. hookeriana*) genetic variation results in a more complex plant–insect food web due to complementarity in trophic interactions. Specifically, we found that the average complexity (LD_q , quantitative-weighted linkage density) of the plant–insect food web increased by 20% over the range of genetic variation (number of genotypes) in the experimental population of willows. Gray circles correspond to the average food-web complexity estimates for each replicate simulation ($n = 50$ for each level of genetic variation), whereas blue circles correspond to the overall average complexity of food webs at each level of genetic variation. Black circles correspond to the average complexity of one-genotype mixtures at four different levels of sampling effort (i.e., number of plants sampled), and the dashed line represents the predicted increase in complexity of one-genotype mixtures with greater sampling effort. The *Inset* shows how the average composition of trophic interactions (willow–gall and gall–parasitoid) differed by 73% among willow genotypes (PERMANOVA on Bray–Curtis dissimilarities, $F_{22,89} = 1.90$, $P = 0.001$), suggesting an important role of complementarity in determining food-web complexity. In this ordination plot, black letters and gray ovals correspond to the centroid and SE of the centroid, respectively, for the composition of trophic interactions found on each willow genotype. Centroids and their SEs were calculated from a constrained analysis of principal coordinates (CAP) on Bray–Curtis dissimilarities.

and food-web complexity, we used a structural equation model (*Materials and Methods*). We found that increasing genetic variation resulted in a more diverse community of galls and a more generalized network of gall–parasitoid interactions, albeit through two main pathways (*SI Appendix, Fig. S2*). On the one hand, increasing genetic variation resulted in higher gall species richness, which had a positive direct effect on food-web complexity (standardized path effect = 0.21). On the other hand, increasing genetic variation resulted in higher gall abundances, which indirectly increased complexity by increasing the effective number of parasitoid species per gall (standardized path effect = 0.26). Other pathways had comparatively small and idiosyncratic effects on food-web complexity (*SI Appendix, Fig. S2*).

An important limitation of our simulation and experimental design is that we were unable to estimate the extent to which food-web complexity is influenced by nonadditive effects of genetic

variation. Nonadditive effects may arise in a variety of ways (e.g., competition and facilitation, associational resistance/susceptibility, source-sink dynamics), and prior work has shown that host-plant genetic variation can have positive (26), neutral (27), or negative (28) nonadditive effects on the diversity of upper trophic levels. Future experiments are needed that explicitly manipulate levels of genetic variation and test for the presence and magnitude of non-additive effects on food-web structure. It is worth noting, however, that our qualitative conclusion, namely that genetic variation likely increases food-web complexity, will still hold unless negative, nonadditive effects are equal or greater in magnitude compared with the additive effect that we observed.

Conclusions. Our results suggest that the gain or loss of genetic variation within a key species may fundamentally alter food-web complexity and therefore the persistence of food webs. There are two main conclusions from our work. First, intraspecific variation in multiple traits is an important driver of network structure; therefore, mechanistic models of food-web structure should incorporate such variability within species (7), as this can enhance the accuracy of these models in predicting trophic interactions (29). Given that plants, insect herbivores, and their parasitoids comprise over half of all known species of metazoans (30, 31), accounting for intraspecific variation in a wide range of functional traits should be a priority for future food-web models (32). Second, understanding the direct and indirect effects of genetic variation on trophic interactions is essential for predicting how evolutionary processes will affect the structure and persistence of food webs over time. Indeed, our simulations suggest that the loss of genetic variation will result in less complex food webs. Moreover, genetic variation provides the raw material for evolution by natural selection; therefore, losing genetic variation in key species may hinder the adaptive capacity of both the species and the food web under future environmental change (33, 34). At this point, however, we are currently lacking a theoretical and empirical understanding of how genetic variation scales up to affect the dynamics of food webs. Given that the current rate of population extinction is orders of magnitude higher than the rate of species extinction (35), our study highlights the pressing need for research examining how the loss of genetic variation within and among populations will affect food webs and the ecosystem services that they provide (36, 37).

Materials and Methods

Common Garden Experiment and Plant Traits. To isolate the effects of coastal willow (*S. hookeriana* Barratt ex Hooker) genetic variation on the plant-insect food web, we used a common garden experiment consisting of 26 different willow genotypes (13 males; 13 females), located at Humboldt Bay National Wildlife Refuge (HBNWR) (40° 40' 5" N, 124° 12' 4" W) near Loleta, CA. Willow genotypes were collected from a single population of willows growing around Humboldt Bay. Although relatedness among these genotypes is unknown, their phenotypes in multivariate trait space are quite distinct from each other (details in *SI Appendix*), suggesting that we can treat them as independent from one another. This common garden was planted in February 2009 with 25 clonal replicates (i.e., stem cuttings) of each willow genotype in a completely randomized design in 2 ha of a former cattle pasture at HBNWR. Willows in our garden begin flowering in February and reached their peak growth in early August. During this study, willows had reached 2–4 m in height. Further details on the genotyping and planting of the common garden are available in ref. 18.

To identify the plant traits that may be determining resistance to galling insects, we measured 40 different traits associated with leaf quality (36 traits) and plant architecture (4 traits). Each of these 40 traits exhibited significant, broad-sense heritable variation (mean leaf quality $H^2 = 0.72$; mean architecture $H^2 = 0.27$; range of H^2 for all traits = 0.15–0.97). For further details on how these willow traits were sampled and quantified, see methods in ref. 18. We then reduced these 40 traits into 13 composite traits that had a negligible degree of multicollinearity using principal component analysis, sequential regression (residuals of one trait after accounting for correlation between two traits), or removing one trait from a pair of highly correlated traits (details in methods in ref. 18). The final set of leaf quality traits

included salicylates/tannins PC1, flavones/flavonols PC1–2, phenolic acids PC1–2, flavanones/flavanonols PC1 (table S3 of ref. 18), C:N, water content, specific leaf area (residuals from water content), and trichome density. The final set of plant architecture traits included plant size, plant height (residuals from plant size), and foliage density (residuals from plant size).

Quantifying the Genetic Specificity of the Plant–Insect Food Web. To build a quantitative food web for each willow genotype, we collected galls from about five randomly chosen replicates of each genotype in September 2012 ($n = 145$ willows, range: four to nine replicates per genotype). For each replicate willow, we collected all galls occurring on one randomly selected basal branch. We restricted our gall collections to those induced by midges in the insect family Cecidomyiidae (four species). These species included a leaf gall (*Iteomyia saliciverruca*), bud gall (*Rabdophaga salicisbrassicoides*), apical-stem gall (unknown midge species), and midstem gall (*Rabdophaga salicisbattatus*). To quantify the abundance of gall–parasitoid interactions, we placed collected galls into 30-mL plastic transport vials (loosely capped at the end), which we maintained at room temperature in the laboratory for 4 mo. We then opened galls under a dissecting scope and determined whether the gall survived or was parasitized and, if parasitized, the identity of the parasitoid species. In total, we identified five species of hymenopteran parasitoids, including *Platygaster* sp. (Family: Platygasteridae), *Mesopolobus* sp. (Family: Pteromalidae), *Torymus* sp. (Family: Torymidae), *Tetrastichus* sp. (Family: Eulophidae), and an unknown species of Mymaridae (hereafter, Mymarid sp. A), as well as one predatory midge (*Lestodiplosis* sp., Family: Cecidomyiidae). This predatory midge is functionally similar to the other parasitoids so we collectively referred to this natural enemy community (six species) as parasitoids for brevity. All together, we documented 12 unique gall–parasitoid interactions (Fig. 1), which appear to represent the vast majority of interactions in the gall–parasitoid network (details in *SI Appendix*). We omitted from analysis those galls for which we could not reliably determine the cause of mortality. We quantified gall abundance by counting the number of surviving and parasitized larva for each gall species collected from each branch. For gall size, we measured galls to the nearest 0.01 mm at their maximum diameter (perpendicular to the direction of plant tissue growth).

To quantify the genetic specificity of trophic interactions with galling insects, we tested for differences in gall sizes, abundances, and community composition among willow genotypes. For gall size, we analyzed separate linear models with willow genotype as the predictor variable and average gall size as the response variable, but we weighted the analysis by the number of galls used to calculate average gall size. We weighted the analysis because we expected that averages based on more galls reflect a more accurate estimate of the average size of galls found on a willow individual. For gall abundances, we analyzed multivariate generalized linear models (multivariate GLMs, error distribution = negative binomial, link function = log) with willow genotype as the predictor variable and a matrix of gall abundances as the response variable. For gall community composition, we used permutational MANOVA (PERMANOVA) with willow genotype as the predictor variable and a matrix of Bray–Curtis dissimilarities in gall abundances as the response variable. To identify the plant traits mediating resistance to galling insects, we used the same analyses as for gall sizes (weighted linear models) and abundances (multivariate GLMs) except that our predictor variable was now a matrix of willow traits. To select a final model of willow traits, we sequentially removed traits based on Aikake information criteria (AIC) to identify a nested set of candidate statistical models. We then used likelihood ratio tests to identify the statistical model of willow traits that best predicted gall abundances or gall sizes.

To quantify the genetic specificity of the network of gall–parasitoid interactions, we tested for differences in the abundance, composition, and strength of gall–parasitoid interactions among willow genotypes. For the abundance and composition of gall–parasitoid interactions, we used the same analytical approach as we did to test for differences in gall abundances and community composition. For these analyses, however, we had a matrix of the abundance (multivariate GLMs) or dissimilarity (PERMANOVA) of unique gall–parasitoid interactions as the response variable. To identify the mechanisms determining the abundance of gall–parasitoid interactions, we again used multivariate GLMs except that our predictor variable was now a matrix of gall abundances and gall sizes. We then used the same approach as we did to identify the willow traits that best predicted gall abundances (i.e., AIC and likelihood ratio tests) to identify which gall sizes and abundances best predicted the abundance of gall–parasitoid interactions. For the strength of gall–parasitoid interactions, we used separate GLMs (error distribution = binomial, link function = logit) with willow genotype as the predictor variable and the proportion of galls parasitized as our response variable for each gall species. If we detected an effect of willow genotype on total parasitism rates, then we analyzed separate GLMs for each parasitoid species to determine which parasitoids were driving total parasitism rates.

Finally, we again used AIC and likelihood ratio tests to examine whether parasitism rates were due to gall abundance, gall size, or their interaction.

Simulating the Additive Effects of Genetic Variation on Network Complexity.

For our index of complexity, we chose to use quantitative-weighted linkage density, LD_q , which is based on Shannon diversity and is the average of the effective number of prey and predatory interactions for a given species, weighted by their energetic importance (details on how LD_q was calculated are available in *SI Appendix* and in refs. 38 and 39). LD_q (hereafter, food-web complexity) is less sensitive to variation in sample size compared with other measures of food-web complexity (39), making it an appropriate measure of complexity for our study.

To examine whether genetic variation increases food-web complexity, we designed a resampling procedure to estimate the complexity of the plant–insect food web at different levels of genetic variation (range: 1–25 genotype mixtures) from our empirical data. We omitted 1 of the 26 genotypes from this analysis (genotype U) because we did not find any galls on the branches that we sampled. Our resampling procedure consisted of the following two steps. (i) *Generate quantitative matrices*: To ensure willow genotypes had equal sampling effort, we randomly sampled four individual willows of each genotype (without replacement) and their corresponding trophic interactions (willow–gall and gall–parasitoid). Next, we calculated the total abundance of each trophic interaction associated with each genotype, resulting in a quantitative matrix of 25 genotypes (rows) and 16 unique trophic interactions (columns, four willow–gall and 12 gall–parasitoid). (ii) *Sampling genetic variation*: With this matrix, we randomly sampled 1–25 genotypes (without replacement), 200 times each, and calculated the total abundance of each trophic interaction associated with each level of genetic variation. We removed redundant combinations of genotypes that were generated by our random sampling. We then calculated food-web complexity for each sample and then calculated the average complexity for each level of genetic variation. Finally, we repeated this sampling procedure on 50 different matrices to quantify the variability in our estimates of average food-web complexity. This resampling procedure is analogous to methods used in experimental studies (e.g., 26, 27) to estimate the expected additive effects of genetic variation on arthropod diversity.

One constraint of our experimental design and resampling procedure is that estimates of complexity from mixtures with more genotypes are based off more plants (e.g., 1-genotype, 4-plant mixtures vs. 25-genotype, 100-plant mixtures). This would not be a problem if, for example, we had measures of trophic interactions on 25 replicate plants of each willow genotype because we could directly compare 1-genotype, 25-plant mixtures with 25-genotype, 25-plant mixtures. Therefore, it is important to account for the increase in food-web complexity that may come from simply sampling more plants. We estimated this sampling effect by first using our resampling procedure to generate 1,000 estimates of average complexity for one-genotype mixtures based on progressively higher levels of sampling effort (one to four plants). We then used an asymptotic model (40) to predict the average complexity of food webs in 1-genotype, 100-plant mixtures to use as a baseline for estimating the additive effects of genetic variation (Fig. 6, dashed line). Details of the asymptotic model and our evaluation of alternative models are given in *SI Appendix*.

To examine the pathways by which genetic variation influences food-web complexity, we built a piecewise structural equation model (details given in *SI Appendix*) using data from 1 of the 50 replicates of our resampling procedure. We observed the same qualitative results when we explored other replicates, so we report only the quantitative results from the first replicate.

All statistical analyses were conducted in R version 3.1.2 (41).

ACKNOWLEDGMENTS. We thank E. Wu, E. Nelson, the staff of Humboldt Bay National Wildlife Refuge (US Fish and Wildlife Service), and numerous undergraduates from Humboldt State University for helping establish and maintain the willow common garden. L. Mackas-Burns, B. Locke, M. DeSiervo, and J. Jackson provided valuable assistance with the fieldwork and S. Sorsa assisted with leaf phenolic extractions. Comments from M. Rodriguez-Cabal, D. Gravel, and two anonymous reviewers improved the quality of the manuscript substantially. M.A.B. was supported by a Biodiversity Research: Integrative Training and Education Fellowship, a Graduate Student International Research Mobility Award, and a Four-Year Fellowship from the University of British Columbia. J.B. and M.A.F. were supported by a European Research Council Advanced Grant (to J.B.). R.J.-T. was supported by the Academy of Finland (Grant 267360). G.M.C. was supported by the Miller Institute for Basic Research in Science as well as a Natural Sciences and Engineering Research Council Discovery grant.

1. Stouffer DB, Bascompte J (2011) Compartmentalization increases food-web persistence. *Proc Natl Acad Sci USA* 108(9):3648–3652.
2. Rohr RP, Saavedra S, Bascompte J (2014) Ecological networks. On the structural stability of mutualistic systems. *Science* 345(6195):1253–1257.
3. Dunne J, Williams R, Martinez R (2002) Network structure and biodiversity loss in food webs: Robustness increases with connectance. *Ecol Lett* 5(4):558–567.
4. Clark JS (2010) Individuals and the variation needed for high species diversity in forest trees. *Science* 327(5969):1129–1132.
5. Violle C, et al. (2012) The return of the variance: Intraspecific variability in community ecology. *Trends Ecol Evol* 27(4):244–252.
6. Bolnick DI, et al. (2011) Why intraspecific trait variation matters in community ecology. *Trends Ecol Evol* 26(4):183–192.
7. Poisot T, Stouffer DB, Gravel D (2015) Beyond species: Why ecological interaction networks vary through space and time. *Oikos* 124(3):243–251.
8. Bailey JK, Wooley SC, Lindroth RL, Whitham TG (2006) Importance of species interactions to community heritability: A genetic basis to trophic-level interactions. *Ecol Lett* 9(1):78–85.
9. Abdala-Roberts L, Mooney KA (2013) Environmental and plant genetic effects on tri-trophic interactions. *Oikos* 122(8):1157–1166.
10. Fritz RS (1995) Direct and indirect effects of plant genetic variation on enemy impact. *Ecol Entomol* 20(1):18–26.
11. Fritz RS, Price PW (1988) Genetic variation among plants and insect community structure: Willows and sawflies. *Ecology* 69(3):845–856.
12. Maddox GD, Root RB (1990) Structure of the encounter between goldenrod (*Solidago altissima*) and its diverse insect fauna. *Ecology* 71(6):2115–2124.
13. Harmon LJ, et al. (2009) Evolutionary diversification in stickleback affects ecosystem functioning. *Nature* 458(7242):1167–1170.
14. Post DM, Palkovacs EP, Schielke EG, Dodson SI (2008) Intraspecific variation in a predator affects community structure and cascading trophic interactions. *Ecology* 89(7):2019–2032.
15. Bukovinsky T, van Veen FJF, Jongema Y, Dicke M (2008) Direct and indirect effects of resource quality on food web structure. *Science* 319(5864):804–807.
16. Moya-Laraño J (2011) Genetic variation, predator-prey interactions and food web structure. *Philos Trans R Soc Lond B Biol Sci* 366(1569):1425–1437.
17. MacArthur R (1955) Fluctuations of animal populations and a measure of community stability. *Ecology* 36(3):533–536.
18. Barbour MA, et al. (2015) Multiple plant traits shape the genetic basis of herbivore community assembly. *Funct Ecol* 29(8):995–1006.
19. Hawkins BA, Cornell HV, Hochberg ME (1997) Predators, parasitoids, and pathogens as mortality agents in phytophagous insect populations. *Ecology* 78(7):2145–2152.
20. Abrahamson WG, Weis AE (1997) *Evolutionary Ecology Across Three Trophic Levels: Goldenrods, Gallmakers, and Natural Enemies* (Princeton University Press, Princeton, NJ).
21. Whitham TG, et al. (2012) Community specificity: Life and afterlife effects of genes. *Trends Plant Sci* 17(5):271–281.
22. Yoshida T, Jones LE, Ellner SP, Fussmann GF, Hairston NG, Jr (2003) Rapid evolution drives ecological dynamics in a predator-prey system. *Nature* 424(6946):303–306.
23. Agrawal AA, Hastings AP, Johnson MTJ, Maron JL, Salminen J-P (2012) Insect herbivores drive real-time ecological and evolutionary change in plant populations. *Science* 338(6103):113–116.
24. Moya-Laraño J, et al. (2012) Climate change and eco-evolutionary dynamics in food webs. *Adv Ecol Res* 47:1–80.
25. Melián CJ, et al. (2011) Eco-evolutionary dynamics of individual-based food webs. *Adv Ecol Res* 45:225–268.
26. Crutsinger GM, et al. (2006) Plant genetic diversity predicts community structure and governs an ecosystem process. *Science* 313(5789):966–968.
27. Johnson MTJ, Lajeunesse MJ, Agrawal AA (2006) Additive and interactive effects of plant genetic diversity on arthropod communities and plant fitness. *Ecol Lett* 9(1):24–34.
28. McArt SH, Cook-Patton SC, Thaler JS (2012) Relationships between arthropod richness, evenness, and diversity are altered by complementarity among plant genotypes. *Oecologia* 168(4):1013–1021.
29. Woodward G, et al. (2010) Individual-based food webs: Species identity, body size and sampling effects. *Adv Ecol Res* 43:211–266.
30. Price PW (1980) *Evolutionary Biology of Parasites* (Princeton University Press, Princeton, NJ).
31. Strong DR, Lawton JH, Southwood SR (1984) *Insects on Plants: Community Patterns and Mechanisms* (Harvard University Press, Cambridge, MA).
32. Henri DC, van Veen FJF (2011) Body size, life history and the structure of host–parasitoid networks. *Adv Ecol Res* 45:135–180.
33. Carroll SP, et al. (2014) Applying evolutionary biology to address global challenges. *Science* 346(6207):1245–1249.
34. Jump AS, Marchant R, Peñuelas J (2009) Environmental change and the option value of genetic diversity. *Trends Plant Sci* 14(1):51–58.
35. Hughes JB, Daily GC, Ehrlich PR (1997) Population diversity: Its extent and extinction. *Science* 278(5338):689–692.
36. Luck GW, Daily GC, Ehrlich PR (2003) Population diversity and ecosystem services. *Trends Ecol Evol* 18(7):331–336.
37. Schindler DE, et al. (2010) Population diversity and the portfolio effect in an exploited species. *Nature* 465(7298):609–612.
38. Bersier L-F, Banašek-Richter C, Cattin M-F (2002) Quantitative descriptors of food-web matrices. *Ecology* 83(9):2394–2407.
39. Banašek-Richter C, et al. (2009) Complexity in quantitative food webs. *Ecology* 90(6):1470–1477.
40. Colwell RK, Coddington JA (1994) Estimating terrestrial biodiversity through extrapolation. *Philos Trans R Soc Lond B Biol Sci* 345(1311):101–118.
41. R Core Team (2014) R: A Language and Environment for Statistical Computing (R Foundation for Statistical Computing, Vienna).